

Effect of noise for two interacting particles in a random potential

F. BORGONOV ^[a]

Dipartimento di Matematica, Università Cattolica, via Trieste 17, 25121 Brescia, ITALY

D.L. SHEPELYANSKY ^[b]

Laboratoire de Physique Quantique, Université Paul Sabatier,

118, route de Narbonne, 31062 Toulouse Cedex, FRANCE

Abstract: We investigated the effect of noise on propagation of two interacting particles pairs in a quasi one-dimensional random potential. It is shown that pair diffusion is strongly enhanced by short range interaction comparing with the non-interacting case.

PACS numbers 71.55.Jv, 72.10Bg, 05.45.+b

Recent investigations showed that two interacting particles (TIP) in a random potential can propagate coherently on a distance l_c which is much bigger than the one-particle localization length l_1 without interaction [1, 2, 3, 4, 5, 6]. According to [1, 2] the TIP localization length is given by:

$$l_c \sim l_1^2 M (U/V)^2 / 32 \quad (1)$$

where M is the number of transverse channels in a quasi one-dimensional wire, U is the strength of on-site interaction assumed to be comparable to or less than the bandwidth and V is the hopping matrix element between nearby sites. Here it is also assumed that the wave vector k_F corresponding to TIP energy is $k_F \sim 1/a = 1$, a being the lattice constant. While the exact verification of (1) is still under investigation the existence of the enhancement of two particle localization length have been clearly demonstrated in numerical simulations [3, 4, 5, 6]. These simulations have been done for different models. Main results are for 1-d Anderson model with TIP [3, 5] and for the model of two interacting kicked rotators [4]. This last model is very convenient for the investigation of wave packets spreading in time due to the existing effective numerical methods. For this reason, in our numerical studies, we used the last model.

The problem we want to address in this letter is the influence of noise on TIP localization. For one particle the effect of noise has been analyzed during the last years and the physics of this phenomenon is well understood [7, 8, 9, 10]. Generally, the main effect of noise is the destruction of interference after a coherence time t_c , after which the particle makes a jump on a distance l_1 that leads to a diffusion rate $D_1 \sim l_1^2/t_c$ along the chain. For high frequency noise with amplitude ϵ this time is $t_c \propto 1/\epsilon^2$ [7]. In the case of low

frequency (ω) noise $t_c \propto (\epsilon\omega)^{2/3}$ [10]. Here we only analyzed the case of high frequency noise on TIP problem.

We studied the model of two interacting kicked rotators in presence of noise. The evolution operator is given by :

$$\hat{S} = \exp\{-i[H_0(\hat{n}_1) + H_0(\hat{n}_2) + U\delta_{n_1,n_2}]\} \times \exp\{-i[V(\theta_1, t) + V(\theta_2, t)]\} \quad (2)$$

where $\hat{n}_{1,2} = -i\partial/\partial\theta_{1,2}$, $H_0(\hat{n}) = T\hat{n}^2/2$, $V(\theta, t) = [k + \epsilon f(t)]\cos\theta$, and $f(t)$ is a random function of t homogeneously distributed in the interval $[-1, 1]$.

For $U = 0$ and $\epsilon = 0$ we have two decoupled kicked rotators and in the chaos domain $kT > 1$ the localization length is $l_1 \approx k^2/2$ [11]. In the presence of noise $\epsilon > 0$ the decoherence time is $t_c \sim 1/\epsilon^2$. If this time is less than the localization time $t_1^* \approx l_1$ localization effects are not important and diffusion goes with the usual classical rate $D \approx k^2/2$. On the other side when $t_1^* < t_c$ the diffusion rate is $D_1 \sim \epsilon^2 l_1^2$ [7].

In the presence of interaction ($U \neq 0$), but without noise ($\epsilon = 0$), a TIP pair of size l_1 propagates on a distance $l_c \gg l_1$ and it is localized after a time $t_2^* \sim l_c l_1$ [12]. On a time interval $t_1^* < t < t_2^*$ the pair diffuses with a diffusion rate $D_p \sim U^2 D$. The noise leads to a destruction of localization after a decoherence time $t_c \sim 1/\epsilon^2$ as in the one-particle case. After this time the pair makes a jump of size l_c and therefore the noise-induced pair diffusion rate can be estimated as $D_+ \sim l_c^2/t_c$ that gives for $t_c > t_2^*$:

$$D_+ \sim \epsilon^2 l_c^2 \sim D_1 (l_c/l_1)^2 \quad (3)$$

This means that for $l_c/l_1 \gg 1$ the noise-induced TIP diffusion rate is strongly enhanced with respect to the non interacting one (D_1). In the case of relatively strong

noise, $t_c < t_2^*$ and D_+ becomes comparable with the pair diffusion rate D_p on a time scale $t_1^* < t < t_2^*$.

The estimate (3) gives the diffusion rate of the center of mass of the TIP pair. However noise also leads to a separation of two particles. This separation goes in a diffusive way with a diffusion rate D_1 which is independent from interaction U . Due to this on asymptotically large times the pair propagation will be subdiffusive. At this time the spreading of the center of mass Δn_+^2 can be estimated in the following way $\Delta n_+^2 \sim \nu D_+ t$ where ν is the probability of collision between two particles $\nu \sim l_1/\Delta n_-$. Here Δn_- is the effective pair size at the time t which in turn can be estimated as $\Delta n_-^2 \sim D_1 t$. Therefore we have $\Delta n_+^2 \sim (l_c/l_1)^2 l_1 \sqrt{D_1 t}$.

However on short times, when $\Delta n_- \sim l_1$ the diffusive spreading in n_+ will be given by $\Delta n_+^2 \sim D_+ t$. Examples of these diffusive spreading are presented in Fig.1 and Fig.2 where the growth of $\sigma_+ = (|n_1| + |n_2|)^2/4$ and $\sigma_- = (|n_1| - |n_2|)^2$ is shown as a function of time. These results clearly show that the diffusion rate $D_+ = \sigma_+/t$ is strongly enhanced with the interaction switching on (approximately 20 times from $U = 0$ to $U = 2$). At the same time the diffusion rate in n_- ($D_- = \sigma_-/t$) remains practically the same (see Fig.2). For sake of comparison we also present in Figs. 1,2 TIP localization in absence of noise, when $\sigma_{\pm}$ are oscillating in time near their asymptotic values.

To check the relation (3) we determined the diffusion rate D_+ and checked its dependence on parameters. According to (3) $D_+ \sim \epsilon^2 \sigma_+^0$ where $\sigma_+^0 \sim l_c^2$ is the asymptotic value of σ_+ in absence of noise. The dependence of D_+/ϵ^2 on σ_+^0 is shown in Fig. 3 for different ϵ, U and k values. The average behaviour is given by the approximate relation

$D_+ = 13\epsilon^2\sigma_+^0$ and it is in agreement with the theoretical prediction (3).

In real systems, noise can appear as the result of electron interaction with phonons at finite temperature. Our results indicate that the noise produced by phonons can lead to a strong enhancement of diffusion (conductance) of electrons in a random potential. Of course in the analysis of a physical model the case of finite particles (or quasi-particles) density should be considered. In this case the probability to find two particles within a distance l_1 one from each other is of the order of $\mathcal{W} \sim l_1\rho$ where ρ is the linear (per unit length) density of particles. This leads to a decrease of the effective diffusion rate which in this case can be estimated as $D_{eff} \sim \mathcal{W}D_+ \sim D_1 l_c^2 \rho / l_1$. Even for small density D_{eff} can be larger than D_1 and we expect that the effect of interaction-enhanced diffusion is physically relevant. However future investigations on the finite particles density case should be done.

We acknowledge the University of Como for hospitality during the final stage of this work.

References

- [a] Also Istituto Nazionale di Fisica della Materia , Unità di Milano, via Celoria 16,
22100 Milano, Italy
and Istituto Nazionale di Fisica Nucleare, sezione di Pavia, via Bassi 7, 27100
Pavia, Italy
- [b] Also Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia
- [1] D.L.Shepelyansky, Phys. Rev. Lett. **73**, 2607 (1994).
- [2] Y.Imry, Europhys. Lett. **30**, 405 (1995).
- [3] K. Frahm, A. Müller-Groeling, J.-L. Pichard and D. Weinmann, Europhys. Lett.
31, 405 (1995); D. Weinmann, A. Müller-Groeling, J.-L. Pichard and K. Frahm,
Phys. Rev. Lett. **75**, 1598 (1995).
- [4] a) F. Borgonovi and D.L. Shepelyansky, Nonlinearity **8**, 877 (1995); b) J. de Phys. I
France **6**, 287 (1996).
- [5] F. von Oppen, T. Wettig and J. Müller, Phys. Rev. Lett. **76**, 491 (1996), F. von Oppen
and T. Wettig, Europhys. Lett. **32**, 741 (1995)
- [6] K. Frahm ,A. Müller-Groeling and J.L.Pichard, Phys. Rev. Lett. **76**, 1509 (1996).
- [7] E.Ott, T.M. Antonsen Jr and J.D.Hanson, Phys. Rev. Lett. **53**, 2187 (1984).
- [8] D.Cohen, Phys. Rev. A **44**, 2292 (1991).
- [9] S.Fishman and D.L.Shepelyansky, Europhys. Lett. **16**, 643 (1991).
- [10] F.Borgonovi and D.L.Shepelyansky Phys. Rev. E **51**, 1026 (1995).

- [11] D.L.Shepelyansky Physica D **28**, 103 (1987).
- [12] D.L.Shepelyansky, Proceeding of " Recontres de Moriond" (1996) in press.

Figure captions

Fig. 1: Dependence of σ_+ on time for different U values: $U = 0, 1, 2$ correspond to lower, middle and upper full curves, $\epsilon = 0.02, k = 4, kT = 5$. Dotted line is for $\epsilon = 0, U = 2, k = 4, kT = 5$. Initially particles are at $n_1 = n_2 = 0$. The basis is $-250 < n_{1,2} < 250$.

Fig. 2: The same as Fig.1 but for σ_- .

Fig. 3: Dependence of TIP pair diffusion rate on σ_+^0 . Full circles are for $k = 4, \epsilon = 0.01, 0 \leq U \leq 2$, open circles for $k = 4, \epsilon = 0.02, 0 \leq U \leq 2$, full squares for $k = 5.7, \epsilon = 0.01, U = 0, 1$, open squares for $k = 4.8, \epsilon = 0.02, U = 1, 2$, up-triangles for $k = 4.8, \epsilon = 0.01, U = 1, 2$, down-triangles for $k = 3.3, \epsilon = 0.01, U = 1, 2$, diamonds for $k = 3.3, \epsilon = 0.03, U = 1, 2$, asterisks for $k = 4, \epsilon = 0.05, U = 0, 1$. In all cases the chaos parameter has been fixed $kT = 5$. Dashed line shows average dependence $D_+ = 13\epsilon^2\sigma_+^0$.





